

**COMPOSITIONS COMPRISING GENOME SEGMENTS
AND METHODS OF USING THE SAME**

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Application No.
5 60/174,155 filed January 3, 2000, which is incorporated herein by reference.

FIELD OF THE INVENTION

The present invention relates to compositions and methods of treating individuals with diseases and disorders associated with genetic mutations. According to the present invention, DNA is administered to individuals who take it up and replace DNA
10 with mutated sequences with DNA with non-mutated sequences or substitute existing set of alleles to a different one. The present invention also relates to compositions and methods for preventing and treating infections, allergies, transplant rejections and infertility.

BACKGROUND OF THE INVENTION

15 Every cell of an organism contains genomic DNA that keeps encoded information about all proteins of all cells of the organism and eventually about the whole organism and its development. Initially this genomic DNA is identical in all cells of an organism. But as the organism grows genomic DNA of every cell becomes subjected to

mutational pressure due to environmental factors and mistakes of cellular replication and repair machinery. The known cellular DNA repair mechanisms replace damaged or incorrect DNA bases and rejoin ends of DNA after single-strand and double-strand breaks immediately after mutational events. They can also employ the second DNA chain as a template. When the damage is rather long and affects both DNA chains its repair becomes problematic and the damage results in mutation. The double-strand break repair mechanism can itself be a source of mutations (Pfeifer, P., The mutagenic potential of DNA double-strand break repair. 1998, 96-97, 119-129). As mutations are accumulated in the cellular genome they can cause malfunction of some cells or cancerous transformation.

Environmental mutagenic factors and errors of cellular replication and repair are sources of somatic cell mutations. It has been suggested that gradual accumulation of mutations in cells can cause multi-step cancer transformation. Somatic mutations are considered to be the main cause of aging. Mutations may also induce development of some hereditary associated diseases such as cardio-vascular diseases, high blood pressure, rheumatoid arthritis, and diabetes.

The genetic, multi-mutational nature of cellular cancer transformation and cancer itself suggests that methods of cancer therapy should be directed to the cause of the disease and particularly targeted to the treatment of mutations. Methods for fixing definite point mutations in cells have been suggested (U.S Patent No. 5,795,972 issued to Kmiec on August 18, 1998, which is incorporated herein by reference) but according to the method, mutations must be precisely identified before each treatment. Because not a single mutation has been 100% proven to cause a cancer, the development of these methods requires additional fine and time-consuming research. There is a need for methods which can be applied based upon the current knowledge underlying the causes of cancer.

Applying DNA fragments locally has been suggested for treatment of precancerous conditions in skin of patients and for tanning stimulation (U.S. Patent No. 5,955,059 issued to Gilchrest, et al., on September 21, 1999; and U.S. Patent No. 5,470,577 issued to Gilchrest, et al., November 28, 1995, both of which are incorporated herein by reference). No sequences or sources of DNA are specified. Rather, the patent suggests that 'any appropriate sources' DNA natural or synthetic, for example, salmon DNA with the length from 200 to mononucleotides and nucleosides including dimers, the

09753892.010301

most potent agents in their tests can be used.

Another method to activate DNA repair in cells is offered including delivery of enzymes participating in DNA repair into skin cells (U.S. Patent No. 5,352,458 issued to Yarosh, October 4, 1994; and U.S. Patent No. 5,302,389 issued to Kripke, et. al., April 12, 1994, both of which are incorporated herein by reference). However the latter methods do not suggest fixing of already established or inherited mutations.

There remains a need for methods of correcting genetic mutations. There remains a need for treating individuals who have diseases and disorders associated with genetic mutations. There remains a need to replace undesirable alleles with desirable alleles. There is a further need for methods for inducing tolerance to prevent transplant rejection. There is a further need for methods for inducing tolerance to treat and prevent allergies. There is a need for methods for increasing fertility.

SUMMARY OF THE INVENTION

The present invention arises from the surprising discovery that when an entire genome is administered to an individual in the form of a plurality of polynucleotide molecules that are fragments of genomic DNA, the polynucleotide molecules will circulate, be taken up by cells and translocate to the nucleus of the cells where they will recombine by homologous recombination with the genomic DNA of the cell. Accordingly, genetic defects can be corrected by administering an entire genome to an individual in the form of polynucleotide molecules that are fragments of genomic DNA that are free of the genetic defect. Individuals with diseases and disorders associated with genetic mutations can be effectively treated by the present invention.

Human genomic DNA fragments are used for treating cancer and other diseases caused by mutations. Nucleic acid molecules derived from "healthy" donor DNA is administered to the patient. The nucleic acid molecules correct mutational changes in cells via homologous recombination with mutated genomic sequences. This mechanism modifies allelic genes and in this way an antigenic structure of organism in accordance with composition of administered nucleic acid molecules. The present invention is useful for induction of immunological tolerance at transplantation or pregnancy.

The present invention relates to methods of treating individuals who have

09753892, 010301

diseases or disorders associated with a genetic mutation or undesirable allele in genomic DNA and to methods of preventing individuals from developing diseases or disorders associated with a genetic mutation or undesirable allele in genomic DNA. According to the invention a segment of genomic DNA that has a mutated sequence or undesirable allele

5 is replaced with a corresponding segment of DNA that has a non-mutated sequence or desirable allele. The methods comprise the step of administering to the individual an effective amount of a plurality of polynucleotide molecules that are free of vector sequences. The plurality of polynucleotide molecules collectively comprises an essentially complete genome in polynucleotide molecules having about 100-3000 nucleotides, and

10 includes a polynucleotide molecule which comprises a non-mutated sequence or desirable allele corresponding to the genetic mutation or undesirable allele in the genomic DNA of the cell in the individual. At least some of the plurality of polynucleotide molecules including the polynucleotide molecule which comprises the non-mutated sequence or desirable allele are taken up by the cell of the individual which has the genetic mutation or

15 undesirable allele in genomic DNA, transported to the nucleus of the cell, and recombine with the genomic DNA of the cell by homologous recombination. The mutated sequence or undesirable allele of genomic DNA of the cell is replaced by the non-mutated sequence or desirable allele to correct the genetic mutation in the genomic DNA or incorporate the desirable allele into the genomic DNA.

20 The present invention relates to methods of inducing tolerance and preventing transplant rejection in an individual comprising the steps of administering to the individual an effective amount of a plurality of polynucleotide molecules that are free of vector sequences. The plurality of polynucleotide molecules collectively comprises an essentially complete genome of the donor in polynucleotide molecules having about 100-

25 3000 nucleotides. At least some of the plurality of polynucleotide molecules are taken up by the cell of the individual, are transported to the nucleus of the cell, and recombine with the genomic DNA of the cell by homologous recombination. The genomic DNA of the individual is replaced by genomic DNA of the donor and tolerance is induced and transplant rejection is reduced in the individual.

30 The present invention relates to methods of inducing tolerance and treating allergies in an individual comprising the steps of administering to the individual an

09753892.010301

effective amount of a plurality of polynucleotide molecules that are free of vector sequences. The plurality of polynucleotide molecules collectively comprises an essentially complete genome of the donor in polynucleotide molecules having about 100-3000 nucleotides. At least some of the plurality of polynucleotide molecules are taken up by the
5 cell of the individual, are transported to the nucleus of the cell, and recombine with the genomic DNA of the cell by homologous recombination. The genomic DNA of the individual is replaced by genomic DNA of the donor and tolerance is induced and allergies are reduced in the individual.

The present invention further relates to methods of increasing fertility in a
10 woman. The methods comprise the step of administering to the woman an effective amount of a plurality of polynucleotide molecules that are free of vector sequences. The plurality of polynucleotide molecules collectively comprises an essentially complete genome of a prospective father in polynucleotide molecules having about 100-3000 nucleotides. At least some of the plurality of polynucleotide molecules are taken up by
15 cells of the woman, transported to the nucleus of the cell, and recombine with the genomic DNA of the cell by homologous recombination. Genomic DNA of the woman is replaced by genomic DNA of the prospective father and fertility is improved in the woman.

The present invention relates to methods of preventing or reducing graft versus host disease in a recipient comprising the steps of administering to the donor an
20 effective amount of a plurality of polynucleotide molecules that are free of vector sequences. The plurality of polynucleotide molecules collectively comprises an essentially complete genome of the recipient in polynucleotide molecules having about 100-3000 nucleotides. At least some of the plurality of polynucleotide molecules are taken up by the cells of the recipient, transported to the nucleus of the cell, and recombine with the
25 genomic DNA of the cell by homologous recombination. The genomic DNA of the donor is replaced by genomic DNA of the recipient. Following transplantation/grafting, the level of graft versus host disease is reduced or prevented.

The present invention relates to methods of introducing one or more desirable alleles into livestock which contains undesirable alleles comprising the steps of
30 administering to the livestock an effective amount of a plurality of polynucleotide molecules that are free of vector sequences. The plurality of polynucleotide molecules

09753892.010301

collectively comprises an essentially complete genome of the livestock species in polynucleotide molecules having about 100-3000 nucleotides including the desirable alleles. At least some of the plurality of polynucleotide molecules are taken up by the cells animal, transported to the nucleus of the cell, and recombine with the genomic DNA of the
5 cell by homologous recombination. The undesirable alleles in the genomic DNA of the animal is replaced by the desirable alleles.

The present invention relates to an apparatus for doing large scale PCR preparations. The apparatus comprises a reaction tube that has an inner diameter of 3 mm or more, at least one pump for continuously supplying of reagents to the reaction tube, four
10 temperature chambers and a collection vessel. The reagents are combined and enter the reaction tube by action of the pump. The reaction tube comprises a series of reaction tube lengths which alternately pass through each of the four temperature chambers to produce a cycle segment, such that a length of the reaction tube passes through the first temperature chamber, a length of the reaction tube passes through the second temperature chamber, a
15 length of the reaction tube passes through the third first temperature chamber, a length of the reaction tube passes through the fourth temperature chamber to produce a cycle segment. The reaction tube comprises at least twenty consecutive cycle segments and is connected to the collection vessel following the last cycle segment.

The present invention relates to pharmaceutical compositions that
20 comprises, in a pharmaceutically acceptable carrier, a plurality of polynucleotide molecules which collectively comprise an essentially complete genome in polynucleotide molecules having about 100-3000 nucleotides.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A shows changes in specific protein expression in human cancer
25 cells induced by preparations of human DNA. Three assays were run on each of three cancer cell types (human breast cancer cells BT474, ovarian carcinoma cells OVCAR5 and promyelocytic leukemia cells HL60) and expression was tested of proteins associated with cancer, erbB2 for BT474 and OVCAR5 cells, c-myc for HL60 cells and cyclin D1 for all three cell types. Figure 1A contains six autoradiograms; in each lane C designates the
30 control human breast cancer cells BT474, ovarian carcinoma cells OVCAR5 and

09753892.010301

promyelocytic leukemia cells HL60; each lane D designates the same cells treated for 2 weeks with DNA preparations (2 μ g/well); each lane Ch designates the same cells treated for 2 weeks with chromatin preparations (2 μ g of DNA per well). Equal protein amounts of cell lysates were electrophoresed in SDS-PAAG and transferred to nitrocellulose

5 membranes by Western blotting procedure. The membranes were incubated with specific antibodies to proteins Erb-B2, C-Myc and Cyclin D1, washed in buffer and incubated with secondary antibodies conjugated with HRP according to protocols of Santa Cruz Biotech. Co. The membranes were then washed and developed with Luminol reagent and exposed to X-ray film. Fig.1A shows specific protein expression, X-ray films

10 Figure 1B is a bar graph depicting protein quantitation data corresponding to the data in Figure 1A.

Figure 2 shows the survival of A/He mouse strain specific tumor allograft in immunologically incompatible CBA mice injected with A/He strain DNA preparations. Two types of preparations were used: 1) A/He purified total chromatin digested by
15 micrococcal nuclease to oligonucleosome size fragments (200-3000 bp DNA) (diamond cross, xxxx). 2) A/He isolated genomic DNA mildly sonicated to obtain 200-3000 bp size fragments (///). The following schemes of DNA treatments were applied to 4 groups of the recipient CBA mice: 1 - injection of physiological salt solution (control group); 2 -
20 single i.p. injection of either DNA or chromatin preparation (40 μ g of DNA per injection in the form DNA and chromatin) on the second day after birth; 3 - a series of 4 injections of either DNA or chromatin preparation (40 μ g of DNA per injection in the form of DNA or chromatin) every 5 days starting with the second day after birth; 4 - a series of 3
injections of chromatin (40 μ g of DNA per injection) every 5 days starting from the 6th day after birth. Hepatoma A tumors were inoculated into treated and control mice 25 days
25 after single preparation injection or one week after last injection at multi-injectional therapy. In two weeks approximately 30% of mice that received DNA preparations developed cancer at the spot of inoculation.

Figure 3 shows data from random human genomic DNA amplification in PCR reaction. A random primer PCR procedure was used according to Telenius et al.
30 (1992) with modifications. Telenius's 18-mer partially degenerate oligonucleotide primer DOP 6-MW (5'CCGACTCGAGNNNNNNATGTGG SEQ ID NO:1) was used for random

09753892.010301

DNA amplification in Idaho Technology rapid thermal cycler. Reaction mixture 50 µl contained 1 ng of human DNA, PCR buffer with 0.2 mM MgCl₂, 0.2 mM NTP, 2 µM DOP 6-MW and 40 U/ml Taq polymerase. After denaturing of the mixture at 95°C for 5 min it was incubated 10 sec at 94°C, 10 sec at 53°C and 30 sec at 72°C for different number of times with final exposure 10 min at 72°C. Samples were resolved in 1% agarose gel in TBE buffer and bands were visualized in UV light after ethidium bromide staining. Lanes are corresponding to: 1 to 15 thermal cycles, 2 – 25 cycles, 3 – 35 cycles, 4 – 45 cycles and 5 – molecular weight standards. From the picture one can see that although 15 cycles give low yield, other variants give nearly the same yields of DNA product. Thus using 1 ng of the DNA in a 50 µl reaction mixture we can get up to 10 µg DNA fragments 200-3000 bp in length.

Figure 4 is a schematic drawing of the apparatus for large scale amplification of genomic DNA as described in Example 4.

DESCRIPTION OF PREFERRED EMBODIMENTS OF THE INVENTION

The mechanisms of entering of oligo- and polynucleotides into living organisms, their distribution between tissues and organs, and particularly the methods of their penetration into living cells have been studied. Specific cell surface proteins have been found to bind to nucleic acid molecules and deliver them into the acidic vesicular endosomal compartment inside cells and further on into cell nuclei. Oligonucleotides and polynucleotides have been shown to enter cells via a saturable, partially reversible receptor-mediated endocytosis mechanism. It has been observed that nucleic acids can enter a mammalian organism via gastrointestinal and other mucosas or even through the skin. Having been injected intravenously or intraperitoneally they were delivered systemically by circulation to all cells of the organism.

An even more exciting discovery about these receptors has now been made. The same proteins that bind oligonucleotides on the cell surface are responsible for their delivery into the cell nucleus. Thus nuclear oligonucleotide delivery was found to be a multi-step process employing the cell surface receptors and resulting in reversible redistribution of the receptors in the cell to mostly nuclear localization and the process was activated by extracellular oligonucleotides.

Nucleic acid fragments are normally present in plasma and tissue fluids. They are constantly produced in organisms as a result of cellular death via apoptosis (programmed cell death) and necrosis of cells and consist mainly of the fragmented genomic DNA complexed with nuclear proteins. This fragmented genomic DNA can be taken up by cells of the body, then transported to cellular nuclei and finally can be employed as an external DNA standard for repair of mutations in cells via homologous recombination of the delivered DNA fragments with cellular genomic DNA.

Although the present invention is expressly not intended to be limited by any theory, the data suggest a possible reason related to the discoveries underlying the present invention. The data suggest that there exists a global genomic DNA repair system which uses extracellular DNA fragments from blood circulation or surrounding tissue fluids as a standard for genomic DNA correction in cells by means of homologous recombination, and which works continuously in every cell of the human body although it is unclear yet to what extent this mechanism resists mutational changes of the cellular genome. The global genomic repair system appears to have three main parts or steps of action: i) generation of extracellular fragmented genomic DNA in an organism, ii) cellular uptake and nuclear delivery of the nucleic acid fragments, and iii) homologous recombination of the fragments with genomic DNA. The system stands guard over invariability of the genome and eliminates mutations in every cell of an organism.

Although extracellular nucleic acids can have some misspelled sequences from the death of rare mutated but unrepaired cells, the percentage of such sequences compared to the 'correct' DNA is usually extremely low. However if mutated cells are numerous, they themselves may become a source of mutated plasma DNA (especially if these cells undergo increased apoptosis or necrosis). This can result in multiple mutations of cells in the body. The quantity of mutations will depend on the percentage of mutant DNA in the plasma DNA reservoir. The death of new mutated cells will produce mutated DNA that will increase the proportion of mutated DNA in plasma DNA. This in turn will increase the number of mutated cells via recombination with cellular DNA. An increase in mutated sequences in plasma DNA will be followed by an increase in the proportion of mutated cells in the organism and may further lead to multiple mutations in cells of the body. Depending on how a mutation affects certain properties of cells (e.g. rate of

multiplication or tendency towards apoptosis or necrosis) the percentage of the mutated DNA in plasma DNA will either increase or decrease. Since many different mutations can be displayed in cellular populations of an organism, the mutated sequences will be presented in extracellular DNA of circulation and tissue fluids in corresponding proportions with corrections to the relative rates of cell death in mutated cells. Because such DNA fragments will simultaneously perform repair of mutations in some cells and induce mutations in others, all cells of the organism will be in a state of mobile mutational equilibrium. This equilibrium can be shifted by a variety of factors, such as the possibility for free DNA of the cellular surroundings to play a more important role in the recombination than that of plasma DNA. These processes can change frequencies of mutations in the body and cellular communities and cause gradual drift of the plasma DNA standard. Therefore in light of this theory, cancer and aging may not be diseases of cells or cellular clones, but those of the genome of the organism. If such a global DNA proofreading mechanism exists, any damage or malfunction of its components will result in genetic instability of cells in the organism, accelerating carcinogenesis, premature aging or other disorders.

The theory considered above suggests, that horizontal transfer of genetic material between cells in the body is usually normal state of cellular community. This concerns not only proper inherited genomic DNA but any DNA that happen to be introduced into the body. This DNA will certainly recombine with genomic DNA at sites of homology, and the frequencies (and number of sequences or genes ready to recombine) of recombination will correlate primarily with extent of the relation: high and very high frequencies between DNAs of the same species, lower between remote species and rare between far remote species. Herewith frequencies of recombination for separate homologous genes will be the same regardless of hereditary relations.

Definitions

As used herein, the term "individual" refers to the vertebrate targeted for use of the present invention. Examples of "individuals" contemplated by the present invention include but are not limited to humans, higher order primates, canines, felines, bovines, equines, ovines, porcines, avians, and other mammals.

As used herein, the term "diseases and disorders associated with genetic

mutations" refers to: cancer; heart and blood vessel diseases; peripheral blood vessel diseases; autoimmune diseases; diabetic conditions; neurodegenerative conditions; gastroenterological and hepatological diseases; mutagenic pathogen disorders; classic hereditary diseases; disorders due to exposure to mutagenic stimuli; other multifactorial diseases; hereditary predisposition to one of the above-mentioned diseases; and aging

As used herein, the term "administration" refers to the delivery of polynucleotide molecules to the individual. The term includes, but is not limited to delivery routes including intramuscularly, intravenously, intranasally, intraperitoneally, intradermally, intrathecally, intraventricularly, subcutaneously, transdermally or topically or by lavage. Modes of administration contemplated by this invention include but are not limited to the use of a syringe, intravenous line, transdermal patch, or needleless injector.

As used herein, the term "segment" refers to a portion of genomic DNA with or without mutations.

As used herein, the term "allele" used in the context of an undesirable allele being replaced with a desired allele refers to the entire gene or a portion of the gene such that following recombination with a portion of the gene, the undesirable allele is replaced with the desirable allele. Accordingly, the polynucleotide molecules used to replace an undesirable allele need not comprise the entire gene but enough of the gene to effectively convert the undesirable allele into a more desirable form.

As used herein, the phrase "plurality of polynucleotide molecules that are free of vector sequences" refers to a composition that comprises multiple different nucleic acid molecules which are fragments or copies of fragments of genomic DNA. The nucleic acid molecules are preferably DNA. The nucleic acid molecules do not include sequences other than genomic sequences. A "plurality of polynucleotide molecules that are free of vector sequences" does not contain any vector sequences such as viral or cell sequences from a vector host.

As used herein, the phrase "each polynucleotide molecule has 100-3000 nucleotides" refers to the number of nucleotide bases on a single strand of a nucleic acid molecule. A double stranded nucleic acid molecule such as a DNA would contain 200-6000 nucleotides as 100-3000 base pairs.

As used herein, the phrase "plurality of polynucleotide molecules

09753892.010301

collectively comprises an essentially complete genome" refers to the composition, which if assembled would equal a complete genome. The plurality of polynucleotide molecules are a plurality of fragments of genomic DNA or a plurality of copies of fragments of genomic DNA. For example, in some embodiments, a complete genome of genomic DNA is

5 isolated from an individual, such as isolated chromosomal DNA, fragmented into a plurality of fragments by mechanical sheering and/or sonication and/or partial enzymatic digestion and/or restriction endonuclease digestion. Such a fragmented chromosomal isolate collectively comprises an essentially complete genome. Likewise, copies of an

10 primers and whole genome chromosomal DNA as a template. The plurality of polynucleotide molecules generated by such amplification techniques "collectively comprises an essentially complete genome" in that essentially the entire genomic DNA is copied and found within any one or more of the several randomly produced amplification products which are copies of fragments of genomic DNA.

15 As used herein, the phrase "plurality of polynucleotide molecules includes a polynucleotide molecule which comprises a non-mutated sequence corresponding to the genetic mutation in the genomic DNA of the cell in the individual" is meant to refer to the use of genomic DNA chromosomal starting material from sources believed to be free of the mutated sequence which means that the mutated sequence may be present in such low

20 percentage that in a healthy person it does not induce any pathology. For example, autologous starting material derived from a patient at an early age or prior to exposure to mutagenic stimuli may be used. Similarly, starting material derived from a donor who is not believed to be a carrier or otherwise possess the genetic mutation associated with a patient or at least is not suffering from the disease. In some cases, the exact nature of

25 genetic mutation associated with a patient may be unknown and the starting material derived from a donor who is not patient and therefore believed to be free of the genetic mutation. In some cases, the exact nature of genetic mutation associated with a patient may be known and associated with specific cells, such as tumor cells. The starting material may be derived from non-tumor tissue of the patient and therefore believed to be

30 free of the genetic mutation but this may not be the best idea because even non-tumor tissues of the patient may have the predisposition to develop into a tumor.

09753892.010301

As used herein, the term "treatment" is meant to refer to stabilization, reduction, or reversal of symptoms or the prevention of further advance of disease, debilitation and death.

Diseases and disorders

5 Individuals who have diseases and disorders associated with genetic mutations may be treated by the methods of the invention. As noted above, the diseases include: cancer; heart and blood vessel diseases; peripheral blood vessel diseases; autoimmune diseases; diabetic conditions; neurodegenerative conditions; gastroenterological and hepatological diseases; mutagenic pathogen disorders; classic
10 hereditary diseases; disorders due to exposure to mutagenic stimuli; and other multifactorial diseases.

 The present invention allows for the treatment of multifactorial diseases. With some multifactorial diseases, it is necessary to correct several gene mutations at the same time: for example, higher than normal level of cholesterol is mainly caused by
15 genetic factors. While a few genes have been identified whose mutant alleles have large effects on this trait (e.g. LDLR, familial defective apoB-100), variability in cholesterol levels among individuals in most families is influenced by allelic variation in many genes (polymorphisms) as well as environmental exposures.

 Examples of cancer include: Bladder cancer; brain and CNS cancer; breast
20 cancer; cancer in children; colorectal, esophageal, eye tumors; head and neck; kidney cancer; leukemia; liver cancer; lung cancer; lymphoma; melanoma and other skin cancers; myeloma; pancreatic cancer; prostate cancer; sarcomas; stomach cancer; and testicular cancer. The present invention may be used to prevent metastasis and particularly effective at early stages of cancer, or in some embodiments at pre-cancerous conditions.

25 Examples of heart and blood vessel diseases include: coronary heart disease; angina; heart attack; high blood pressure; congestive heart failure; cardiac arrhythmias; sudden cardiac death; heart valve disease; congenital heart defects; and stroke (usually a consequence of blood vessel disorder);

 Examples of peripheral blood vessel diseases include: peripheral arterial
30 diseases; arteriosclerosis obliterans; Buerger's disease; Raynaud's phenomenon; aneurysms; peripheral venous disorders: varicose veins; and phlebitis (thrombophlebitis).

00753892.010301

Examples of autoimmune diseases includes: Rheumatoid arthritis (RA), multiple sclerosis (MS), Sjogren's syndrome, sarcoidosis, autoimmune thyroiditis, reactive arthritis, ankylosing spondylitis, scleroderma, polymyositis, dermatomyositis, psoriasis, vasculitis, Wegener's granulomatosis, Crohn's disease and ulcerative colitis.

5 Examples of diabetic conditions include Type I and Type II diabetes. In Type I diabetes, also referred to as insulin dependent diabetes mellitus (IDDM), more than 10 different genetic markers have been found to be associated with the disease. In Type II diabetes, also referred to as adult onset or maturity-onset diabetes of the young (MODY) mutations in five different genes, which are code for glucokinase and various transcription
10 factors, have been found to be associated with this form of diabetes.

Examples of neurodegenerative diseases include: Huntington's disease; spinocerebellar ataxias; Alzheimer disease; amyotrophic lateral sclerosis (ALS) and Parkinson's disease.

Examples of gastroenterological and hepatological diseases include:
15 gastrointestinal malignancies, viral hepatitis, hypercholesterolaemia, alpha 1-antitrypsin deficiency; metabolic diseases of the liver and cystic fibrosis.

Examples of mutagenic pathogen disorders include those shown to be caused by nucleic acids of viral and bacterial or fungal pathogens. The severe infections by some pathogens cause mutational changes in the body of the infected individual.
20 Viruses are etiological agents of many human diseases. Viral nucleic acids have a considerable homology with some human genomic sequences. During acute infections large quantities of viral nucleic acids are detected in tissue fluids and circulation. Viral infections are known to induce some mutations in humans, moreover some viruses can incorporate their sequences in human cellular DNA which causes latent, chronic and slow
25 hidden viral infections. Viral infections, such as rubella, result in genetic lesions causing abnormalities of early development. Human genomic DNA preparations can treat post-viral changes in cellular genome. They are likely to treat all forms of viral infections including acute, latent and chronic forms. HIV, herpes simplex virus, rabies virus, influenza virus, hepatitis A, B and C, rubella are all candidates for this treatment.

30 Examples of classic hereditary diseases, i.e. typical human genetic diseases, include, but are not limited to: Achondroplasia, Adult Polycystic Kidney Disorder, Alpha

09753892.010301
T0E0T0.268E5760

Zero Thalassaemia, Alpha₁-Anti-trypsin Deficiency, Alpha-Thalassaemia, Beta-Thalassaemia, Color Vision Disorders, Congenital Hypothyroidism, Congenital Malformations, Cystic Fibrosis, Diabetes Mellitus, Duchenne Muscular Dystrophy, Epilepsy, Familial Hypercholesterolemia (FH), Fetal Rubella Syndrome, Fragile X Mental
5 Retardation, Glucose-6-Phosphate Dehydrogenase Deficiency, Glutathione Deficiency, Haemoglobinopathies, Hemochromatosis, Hemophilia A and B, Huntington Disease, Inherited Thrombophilia, Neurofibromatosis, Obesity, Oculo-cutaneous Albinism, Phenylketonuria (PKU), Sickle Cell Disorders, Thalassemia, and Werding-Hoffman Disorder.

10 Examples of aging as a state of human organism which is characterized by presbyopia, loss of elasticity of skin, wrinkles, hair loss, muscle weakness, fragility of bones, gray hair, flexibility of joints, decline of sexual drive and menopause. One of the main causes of aging is gradual accumulation of mutations in cells of the body. Genomic DNA treatment can prevent and repair mutational changes in cellular genomes. This
15 treatment has a potential to prevent and reverse aging effects in humans.

Examples of disorders due to exposure to mutagenic stimuli include those which result from exposure to ionizing radiation and chemical mutagen exposure.

Examples of other multifactorial diseases includes: dermatological disorders; urological disorders; immunological disorders such as induction of
20 immunological compatibility at transplantations; infertility due to miscarriages; endocrinological disorders; ophtalmological disorders (cataract); otolaryngological disorders; rheumatological disorders; and gynecological disorders.

In some embodiments, the present invention provides the means to correct genetic mutations by replacing mutated sequences with corresponding non-mutated
25 sequences or substitute one set of alleles for a different one. The present invention may be used to replace less desirable alleles with more desirable ones. For example, hair loss can be treated or prevented, hereditary diseases may be considered alleles rather than mutations. Other applications including cosmetic applications may also be allele-based rather than correcting mutations.

30 The methods of the invention can be applied to animal breeding. DNA preparations of animals with successful allelic composition can be prepared and amplified

using this method. When injected in newborn animals or in embryos at early stages of development, these sequences can help accelerate breeding animals with the best properties.

In some embodiments, the present invention provides the means to induce tolerance, reduce transplant rejection and/or graft versus host disease and improve fertility by increasing tolerance/reducing rejection in the mother. In the classical work in early 1950s reported in Billingham et al. (1953) Nature 172(4379):603-606, Medawar and his colleagues found a way to induce artificial tolerance in neonatal mice at allotransplantation via injection of blood lymphocytes from donor to the recipient mouse before transplantation. Later it was found that injections of large amounts of donor splenocytes or bone marrow cells can induce tolerance or diminish the immune response at the transplantation in adult mice. It was suggested that high dose of donor antigens injected caused deaths of lymphocyte clones responsible for the transplant rejection, but a finding that the induced tolerance was unexplainably inheritable complicated the explanation. Recently it was suggested that immune system responds by immunization or tolerance depending of the accompanying presence of some "Danger" signal performed by specific cells (Ridge et al. 1996 Science 271(5256)1723-6).

Immunologists consider injections of cells to recipient just as injection of some quantity of foreign antigen with following reaction of immune system, while every lymphocyte or splenocyte carries set of genomic DNA, and the real sense of the injection may be just injection of foreign nucleic acids into the body. In the light of the theory disclosed in this application for example, the DNA from injected cells is digested partially, taken up by cells of the body and recombines in areas of homology with host DNA. This can modify some genes in cells of the body, including cells of the immune system, and that will change the cells antigenic properties. After this, the immune system will have no choice but to consider its own cells with changed antigenic structure as proper cells of the organism. Lymphocyte clones responsible for recognition of the new antigen are probably deleted. This new property of the body will certainly have the possibility to be inherited if corresponding genes will be modified at least in part of germ cells.

For better results at allotransplantation physicians use lymphocytes or bone marrow cells injections from the future donor, although sometimes this procedure gives

adverse reaction accelerating rejection. When a donor is not known, multiple blood infusions from different persons are administered, to increase allograft survival and diminish immune reaction, although this procedure can allergize the recipient and result in an adverse effect.

5 Instead of whole cells, according to the present invention, either segments of donor DNA or mixed DNA segments of 10-20 healthy persons are administered. These DNA preparations are neither immunogenic, no toxic and will increase tolerance to allograft.

 At xenotransplantation of organs, for example pig's heart to a man, good
10 results are observed when intravenous injections of pig bone marrow cells to the future transplant recipient precede the transplantation procedure. Adverse reaction sometimes results from such infusions. Using pig "DNA" will help to solve this problem and increase xenograft survival. Combined human or recipient DNA injected to a pig donor will minimize or abrogate the graft-versus-host reaction.

15 According to the invention, induction of tolerance can be used to induce tolerance to allergens in those individuals who have allergies. The administration of donor DNA to an individual who has one or more allergies can introduce alleles that prevent allergic reactions when the individual is exposed to the allergen.

 Tolerance of mother to fetus may also be mediated by genetic
20 transformation of a mother's cells with DNA segments of a father's genome originating from sperm ejaculated during sex. Immunologists underline that effect in creation of artificial tolerance by cells injections correlate with number of injected cells to recipient. Some infertility problems or unexpected spontaneous termination of pregnancy can be caused by lack or low level of tolerance created in mother to the fetus because of
25 insufficient quantity of sperm cells injected to women, or problems with possible improper sperm cells degradation, breach of transportation of the DNA segments to circulation, and thus insufficient transformation of mother's cells with the delivered DNA. Such problems may be solved by treating the future mother with prepared segments of the father's DNA, or collected frozen sperm samples of the father, or , if the latter is not possible for some
30 reason, even injections of combined human "DNA" fragments from one or more donors.

Dosage

09753892 "010301
FOEOT" 2682760

The dosage administered varies depending upon factors such as: pharmacodynamic characteristics; its mode and route of administration; age, health, and weight of the recipient; nature and extent of symptoms; kind of concurrent treatment; and frequency of treatment.

- 5 According to some embodiments of the present invention, the plurality of polynucleotide molecules are administered in an amount based on concentrations of circulating DNA and its half life. According to some embodiments of the present invention, the plurality of polynucleotide molecules are administered in an amount of 10 g per month. DNA in blood of a healthy individual approx. 2-10 ng/ml. That is, the
- 10 "normal" concentration of circulating DNA in blood is approximately 2-10 ng/ml. According to preferred embodiments of the invention, polynucleotide compositions are administered in at least a quantity sufficient to increase the normal level by approximately 10X. Depending upon the route of administration, the increase in concentration may be a 10X surge if a complete dose is administered in a short period of time, 5-10 minutes, or
- 15 elevated concentration may be maintained over a longer period of time if the DNA is infused continuously or for a period greater than about ten minutes.

In some embodiments of the present invention, DNA is administered to maintain the concentration in the range of 0.01 – 6 µg/ml, which corresponds to 0.01 – 2 µM concentration of a fragment with average length of 500 b.p.

- 20 Depending upon the manner in which the plurality of polynucleotides are produced, the size distribution will vary. For the purposes of calculating dosage, the fraction of the composition which includes molecules consisting of about 200-3000 nucleotides is measured. Generally, 80% or more of polynucleotides generated by preparations described herein fall within this size distribution. It is preferred that of the
- 25 fraction of the composition which includes molecules consisting of about 200-3000 nucleotides, the average size of about 300-1000, preferably about 500 nucleotides.

- Polynucleotide compositions which comprise 0.4-16 grams of polynucleotide molecules within the 200-3000 nucleotide size distribution are preferably administered to an adult human patient. In some embodiments, the polynucleotide
- 30 compositions comprise 0.8-14 grams of polynucleotide molecules within the 200-3000 nucleotide size distribution are preferably administered to an adult human patient. In some

09753892.010301

embodiments, the polynucleotide compositions comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, or 13 grams of polynucleotide molecules within the 200-3000 nucleotide size distribution are preferably administered to an adult human patient.

In some embodiments, the polynucleotide compositions comprise 0.8-16 grams of polynucleotide molecules within the 300-1000 nucleotide size distribution are preferably administered to an adult human patient. In some embodiments, the polynucleotide compositions comprise 0.8-16 grams of polynucleotide molecules within the 300-700 nucleotide size distribution are preferably administered to an adult human patient.

Since the composition may contain polynucleotides outside the size range, the total amount of polynucleotide may be greater, i.e. up to about 20 grams or more. The dose may be delivered over a period of time, as a divided dose administered in a series of administrations over time or as a divided dose delivered at different sites by multiple routes of administration.

The pharmaceutical compositions according to this aspect of the present invention comprise about 0.4 to 20 g or more of nucleic acid molecules within the 200-3000 size distribution; in some embodiments, about 0.6 to 18 g or more of nucleic acid molecules within the 200-3000 size distribution of DNA; in some embodiments, about 0.8 to 16 g or more of nucleic acid molecules within the 200-3000 size distribution of DNA; in some embodiments, about 1 to 16 g or more of nucleic acid molecules within the 200-3000 size distribution of DNA. In some embodiments of the present invention, the pharmaceutical compositions comprise about 0.4 to 20 g or more of nucleic acid molecules within the 300-2000 size distribution; in some embodiments, about 0.6 to 18 g or more of nucleic acid molecules within the 300-2000 size distribution of DNA; in some embodiments, about 0.8 to 16 g or more of nucleic acid molecules within the 300-2000 size distribution of DNA; in some embodiments, about 1 to 16 g or more of nucleic acid molecules within the 300-2000 size distribution of DNA. In some embodiments, the pharmaceutical compositions according to this aspect of the present invention comprise about 0.4 to 20 g or more of nucleic acid molecules within the 300-1000 size distribution; in some embodiments, about 0.6 to 18 g or more of nucleic acid molecules within the 300-1000 size distribution of DNA; in some embodiments, about 0.8 to 16 g or more of nucleic

09753892.010301

acid molecules within the 300-1000 size distribution of DNA; in some embodiments, about 1 to 16 g or more of nucleic acid molecules within the 300-1000 size distribution of DNA.

In some embodiments, the calculation of DNA dosage is as follows:

5 3 ml of human sperm = about 1 mg human sperm DNA

$$\begin{aligned} & (3 \text{ ml sperm}) \times (70 \times 10^6 \text{ cells/ml}) \times (5 \times 10^{-12} \text{ g [amount of DNA in each spermatozoid]}) \\ & = \text{about } 200 \times 5 \times 10^6 \times 10^{-12} = 1000 \times 10^{-6} \text{ g} = 1 \\ & \text{mg human sperm DNA.} \end{aligned}$$

10 If multiplied by 10,000 (by PCR) = 10 g DNA can be prepared from a 3ml sample.
 $10 \text{ g DNA} / 10 \times 10^{-12} \text{ g DNA/cell} = 10^{12} \text{ cells}$; $10^{12} \times 2000 \mu\text{m}^3$ (volume of embryo fibroblasts) = 2 kg

We have roughly estimated quantities in which cells are distributed in the human organism taking into account the fact that in some tissues there are very few or no

15 cells or there are few cells that actively participate in DNA circulation in the organism.
 The human body of 70 kg equals:

- 1) skin + skin fat - 20 kg (eqv. 0.2 kg cells)
- 2) Bones + connective tissues+ tendons - 30 kg (0.2 kg cells)
- 20 3) Blood - 7 kg (0.07 kg cells)
- 4) Muscles - 10 kg 1/20 (0.5 kg cells)
- 5) Intestine - 3 kg 1/10 (0.3 kg cells)
- 6) liver 1 kg 1/2 (0.5 kg cells)
- 7) Brain 1 kg 1/4 (0.25 kg cells)
- 25 8) Other glands 1 kg (0.25 kg cells)
- Total cells 2.3 kg

Routes of administration

According to the present invention, the polynucleotide molecules may be delivered to the individual by any means and mode of administration that will result in the
 30 polynucleotides being taken up by cells which contain the genetic mutation associated with the disease or disorder to be treated. In some preferred embodiments, the polynucleotide

FOUO: 2685760

molecules are administered by: intravenous injection; intramuscular injection; intradermal injection; subcutaneous delivery; intraperitoneal delivery; topical or lavage delivery by application onto different mucosa and/or skin; ingestion per os (with or without stomach neutralization pretreatment or enteric-coating formulation); per rectum; intravaginally; 5 intraocularly; intranasally, intratumorally; intracerebrally, intraocular injection, by inhalation and by delivery to the spinal cord.

In some applications, particularly cosmetic applications such as for treatment of hair loss or age-damaged and/or sun-damaged skin, topical administration is preferred.

10 The devices and apparatus used can be any of those which are useful to administer the polynucleotide composition by the chosen route of administration. In the case of injections, in addition to syringes, in some embodiments, high powered needleless injection devices or projectile bombardment devices may be used.

Those skilled in the art will perform administration by well known methods 15 including appropriate preparatory and follow up examination.

Regimen

According to the present invention, the polynucleotide molecules may be delivered to the individual by any administration regimen that will maintain sufficient polynucleotide presence such that polynucleotides will be taken up by cells at a level 20 effective to correct genetic mutations. In some preferred embodiments, the polynucleotide molecules are administered continuously. In some embodiments, the polynucleotide molecules are administered hourly. In some embodiments, the polynucleotide molecules are administered daily. In some embodiments, the polynucleotide molecules are administered every other day. In some embodiments, the polynucleotide molecules are 25 administered weekly. In some embodiment, the polynucleotide molecules are administered over the course of 1-26 weeks, preferably 2-13 weeks, preferably 2-8 week, preferably 2-6 weeks and, in some embodiments, more preferably 2-4 weeks..

Formulations

Formulations comprising the polynucleotide molecules are made up 30 according to the mode and site of administration. Such formulation is well within the skill in the art. In addition to nucleic acid molecules, the formulation may also include buffers,

09753892.010301

excipients, stabilizers, carriers and diluents. Suitable pharmaceutical carriers are described in the most recent edition of *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in this field which is incorporated herein by reference.

For parenteral administration, the composition can be, for example,
5 formulated as a solution, suspension, emulsion or lyophilized powder in association with a pharmaceutically acceptable parenteral vehicle. Examples of such vehicles are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Liposomes and nonaqueous vehicles such as fixed oils may also be used. The vehicle or lyophilized powder may contain additives that maintain isotonicity (e.g., sodium chloride, mannitol)
10 and chemical stability (e.g., buffers and preservatives). The formulation is sterilized by commonly used techniques.

The pharmaceutical compositions according to this aspect of the present invention are formulated according to the mode of administration to be used. In cases where injection is the chosen mode of administration, a sterile, isotonic, non-pyrogenic
15 formulation is used. Generally, additives for isotonicity can include sodium chloride, dextrose, mannitol, sorbitol and lactose. Isotonic solutions such as phosphate buffered saline are preferred. Stabilizers include gelatin and albumin.

DNA can be in the form of real solution as sodium or lithium salt or as complexes with basic proteins such as histones or protamines, or synthetic polycations, or
20 precipitates like calcium phosphate, or can be absorbed on carriers such as hydroxylapatite or some anion exchangers. The DNA may be complexed with either synthetic or natural nuclear proteins or both. Further, it may be single- or double-stranded, linear or circular, relaxed or supercoiled.

For administration in the form of suppository DNA can also be
25 formulated as mentioned above (solution, suspension, emulsion, dry powder) in combination with conventionally used inactive base including for instance cocoa butter, starch, methylparaben, propylparaben.

For topical application on skin DNA with commonly used inactive ingredients for lotions, creams and ointments, such as mineral or vegetable oil, propylene
30 Glycol, PEG-Stearate, lecitin, EDTA, methyl/ethylparaben.

For administration in low prophylactic doses DNA solutions in composition

09753892.010301

of lozenges or tablets for dissolving in mouth.

Source

The plurality of polynucleotide molecules may be obtained by processing genomic DNA from a number of sources. In some embodiments, autologous DNA (self donor) that is collected at the patient's young age or at least before the disease or exposure to irradiation or chemical mutagens is used. In such embodiments, genomic DNA may be "banked" from birth or a young age. According to the invention banked collections of DNA samples may be stored in preparation for use according to the invention. In some embodiments, allogenic DNA, such as from living donors or corpses, is used. In such 5
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10140
10145
10150
10155
10160
10165
10170
10175
10180
10185
10190
101

with degenerative primers.

For example, random primer PCR mainly according to Telenius et al (1992) Genomics 13:718-725 which is incorporated herein by reference. Telenius reports 18-mer partially degenerative DOP oligonucleotide primers for random DNA amplification. The typical protocol is as follows: PCR buffer with 0.2 mM MgCl₂, 0.2 mM NTP, 2 μM DOP 6-MW, 40 U/ml Taq polymerase. Cyclor: 95°C 5'; 10" 94°C, 10" 53°C, 30" 72°C (the latter three steps are repeated 30 times), 10' 72°C. Using 1 ng of the DNA in a 50 μl reaction mixture we can get up to 10 μg DNA fragments 200-3000 bp in length. A one liter mixture yields 0.2 g DNA; a 10 liter mixture yields 2 g DNA. For 1 liter mixtures we need 40,000 U Taq polymerase, 100 ml 2 mM NTP (100 mg NTP), 20 ml 100 μM DOP 6 M-W (2 μmol).

Compositions

Another aspect of the invention relates to pharmaceutical compositions that comprise, in a pharmaceutically acceptable carrier a plurality of polynucleotide molecules which collectively comprise an essentially complete genome in polynucleotide molecules having about 100-3000 nucleotides.

In some embodiments, the compositions contain 0.01 – 16 g of polynucleotides having 200-3000 nucleotides each. In some embodiments, the compositions contain 0.01 – 16 g of polynucleotides having 200-3000 nucleotides each with an average length of 300 – 1000 nucleotides. In some embodiments, the compositions contain 0.01 – 16 g of polynucleotides having 200-3000 nucleotides each with an average length of 500 nucleotides. In some embodiments, the compositions contains 0.04 – 16 g of polynucleotides having 200-3000 nucleotides each. In some embodiments, the compositions contain 0.4 – 16 g of polynucleotides having 200-3000 nucleotides each.

In some embodiments, the compositions contain at least 80% of polynucleotide molecules administered are about 200-3000 nucleotides in length. In some embodiments, the compositions contain at least 80% of polynucleotide molecules administered are about 200-3000 nucleotides in length and have an average length of about 300-1000. In some embodiments, the compositions contain at least 80% of polynucleotide molecules administered are about 200-3000 nucleotides in length and have an average length of about 500. In some embodiments, the compositions contain at least 80% of polynucleotide

molecules administered are about 300-2000 nucleotides in length. In some embodiments, the compositions contain at least 80% of polynucleotide molecules administered are about 300-2000 nucleotides in length and have an average length of about 500. In some
5 administered are 300-1000 nucleotides in length. In some embodiments, the compositions contain at least 80% of polynucleotide molecules administered are 300-1000 nucleotides in length and have an average length of 500.

In some embodiments, the pharmaceutical compositions are suitable for injection, i.e. sterile and pyrogen free.

10 EXAMPLE

The foregoing examples are meant to illustrate the invention and are not to be construed to limit the invention in any way. Those skilled in the art will recognize modifications that are within the spirit and scope of the invention. All references cited herein are hereby incorporated by reference in their entirety.

15 Example 1. Preparations of human DNA in cell medium change the pattern of specific protein expression in human cancer cells in culture.

Example 1 demonstrates a capability of DNA preparations prepared from healthy donor sperm cells to influence differential expression of specific cell proteins in human tumor cell lines: breast tumor, BT474, ovarian carcinoma OVCAR5 and promyelocytic
20 leukemia HL60. The breast tumor cells are adhesive epithelial type cells forming a monolayer with multilayer cell clumps; OVCAR5 are fibroblastoid type cells forming a monolayer and HL60 cells are spherical cells that grow in suspension.

Two types of preparations were used:

1) purified total chromatin prepared from sperm cells and fragmented by
25 nuclease digestion to oligonucleosome size (Wagner, et al. Arch. Androl. (1981) 7(3):251-7, which is incorporated herein by reference) (DNA lengths 200-3000 bp).

2) genomic DNA isolated from sperm cells (using Amersham-Pharmacia 'genomic DNA isolation' kit and procedure modified by addition of a reducing agent

09753892, 010301

(DTT) to lysis buffer) mildly sonicated to obtain 200-3000 bp size fragments.

Cells were plated in 24-well plates for tissue culture at a density of 100,000 cells per well and cultured in RPMI 1640 with addition of 5% fetal calf serum and Pen/strep antibiotics at 37°C in the presence of 5% CO₂ in humidified air.

- 5 2 µg of DNA preparations either in the form of DNA solution or as chromatin fragments were added in 20 µl of PBS to all experimental wells, and 20 µl of PBS were added to control wells. The quantity of 2 µg of DNA was chosen in order to provide excess of extracellular DNA over cellular DNA in each well, so that cellular DNA would have a possibility to recombine with extracellular DNA in such manner that all cellular genes
- 10 (cancerous, normal or mutated) could have a good chance to be substituted at least in part with fragments of normal human extracellular DNA. The medium in wells was changed every 3 days and fresh DNA preparations were also added every 3 days for 2 weeks. Following this incubation period, medium was removed and cells were washed with cold PBS 3 times. Cells were then removed from wells, collected by centrifugation and lysed
- 15 in sample loading buffer containing 2% SDS, 20% sucrose, 3mM DTT with incubation of samples on a boiling water bath for 3 min. Samples were resolved in SDS-PAGE and proteins were visualized using an immunoblotting procedure.

- The results are shown in Figures 1A and 1B. Figures 1A and 1B show that the expression levels of epidermal growth factor family protein erbB2, which is overexpressed
- 20 considerably in many different types of cancer, dropped down as a result of DNA treatment in OVCAR5 and BT474 cells. In the cell lines OVCAR5 and HL60 our DNA preparations induced considerable downregulation of protein cyclin D1 whose overexpression in cancer cells often correlates with increased proliferation. In contrast, the treatment does not reduce cyclin D1 levels in BT474 cells and may even cause a slight
- 25 increase. It is noteworthy that BT474 cells are slowly proliferating cells, and may require only a minimal level of cyclin D1 expression that cannot be reduced. The experimental effect appears more pronounced when DNA in complex with basic nuclear proteins (fragmented chromatin) was used as DNA preparation. This can be explained by lesser accessibility of such complexes to nuclease attack in medium or in some cellular
- 30 compartments. It may also may be due to easier cellular uptake of such complexes. It should also be noted that the choice of two weeks time exposure of cells to extracellular

DNA in this experiment was based on preliminary experiments in which we found that treatment with the DNA preparations had no visible toxic effect on cells for at least ten days or more. Furthermore no changes in differential protein expression could be found during first ten days of the experiment. In the following few days (depending on dose) changes in protein expression started to occur, but beyond 20 days, the cells changed their properties so much that it became difficult to maintain cell culture. For example, in BT474 cells the most developed effects were observed in wells with fragmented chromatin; on day 15-16 of the experiment the number of cells in these wells decreased two times compared to the control wells because of the retardation in cell division and death of some cells. Stopping addition of DNA preparations after the 2 weeks' treatment did not help cells detectably: they grew slower and slower. The cells treated with naked DNA also stopped growing, similar to the cells treated with chromatin fragments. We replated these cells, and they attached to plastic, but did not grow well. After a month of such cultivation they were still alive, but did not form a monolayer and their number gradually decreased. This whole pattern looks like the desired solution of many anticancer projects: either make cancer cells die or stop proliferating. Genomic DNA treatment appears to achieve that goal.

**Example 2. Introduction of changes into cellular genomic DNA in living mice.
Induction of artificial tolerance in recipient mice to a donor allograft.**

CBA and A/He are immunologically distinct mouse strains, and A/He allografts to CBA are never accepted but are always quickly rejected (Kaledin, et al. 1974 Conf. of the Res. Lab. of Exp. Biol. Models (Abstracts) Acad. Med. Sci, Moscow USSR p 5-52). The example demonstrates induction of immunological tolerance in CBA mice via injection of appropriate amounts of donor DNA preparations. The supposed mechanism of such induction is that injected donor DNA is substituting responsible alleles in CBA recipient cells to A/He specific alleles. This substitution will result in emergence of A/He antigens on CBA cells, which is probably followed by negative selection of lymphocyte clones normally responsible for the rejection.

A/He mouse strain specific hepatoma A, which is unable to grow in CBA mice mainly because of immunologic incompatibility, was used as tumor allograft for

allotransplantation in treated and control CBA mice. 10^6 tumor cells were inoculated subcutaneously in the lateral part of the hind leg of the recipient CBA mice. The administration of that number of cells results in tumor growth in 100% of the A/He mice over the time of 1 to 2 weeks. This treatment never leads to formation of tumor in CBA mice (Kaledin, et al. 1974 *supra*). DNA isolated from A/He mice liver and mildly sonicated to obtain fragments with lengths between 200 and 3000 bp and A/He mice liver purified chromatin digested with micrococcal nuclease to oligonucleosome size fragments with DNA length between 200 and 3000 bp were injected i.p. in CBA mice according to three different schemes:

- 10 a single i.p. injection of either DNA or chromatin preparation (40 μ g of DNA per injection in the form DNA and chromatin) on the second day after birth;
- a series of 4 injections of either DNA or chromatin preparation (40 μ g of DNA per injection in the form of DNA or chromatin) every 5 days starting with the second day after birth;
- 15 a series of 3 injections of chromatin (40 μ g of DNA per injection) every 5 days starting from the 6th day after birth;
- control groups received injections of physiological salt solution.

Hepatoma A tumors were inoculated into treated and control mice 25 days after a single injection, or one week after the last injection in multi-injectional therapy. Two weeks later, approximately 30% of mice that received DNA preparations developed cancer at the site of inoculation. No cancer developed in control groups of mice. It seemed that the DNA in chromatin fragments was slightly more effective in this experiment. In the following next two weeks, the percentage of mice that developed cancer increased to approximately 50% in treated mice.

- 25 Although immune incompatibility is considered to be the main cause of rejection of hepatoma A tumor by CBA mice, there are probably other factors that interfere with the tumor acceptance. Usually after allotransplantation one can see healing of seams and wounds during the first 10-12 days before the immune response is developed. Then the immune system rejects the allograft. The inoculation of 10^6 of tumor cells to the A/He mouse strain resulted in solid tumor development 7-14 days after the treatment, so if the inoculated tumor cells proliferated at the same rate in CBA mice, then at least a small
- 30

09753892, 010301

tumor should have appeared in the inoculation site before development of the immune response. It did not happen which means that not only immune problems but some other factors or their absence prevented tumor development in the CBA mice soon after the inoculation. If the latter assumption is true then the genomic DNA therapy enables both:

5 creation of the transplantational tolerance and supplying of the CBA mice with some missing, A/He specific factors that are required for hepatoma A proliferation.

Example 3. Random human genomic DNA amplification in PCR reaction.

We used a random primer PCR procedure mainly according to Telenius et al (1992) *supra* with modifications. Telenius's 18-mer partially degenerated DOP 6-MW

10 oligonucleotide primer was used for random DNA amplification. The typical protocol was as follows: PCR buffer with 0.2 mM MgCl₂, 0.2 mM NTP, 2 μM DOP 6-MW, 40 U/ml Taq polymerase. Cyclor: 95°C 5'; 10" 94°C, 10" 53°C, 30" 72°C (the latter three steps are repeated 30 times), 10' 72°C. Using 1 ng of the DNA in a 50 μl reaction mixture we can get up to 10 μM DNA fragments 200-3000 bp in length. 1 liter mixtures yields 0.2 g DNA,

15 10 liter mixtures yields 2 g DNA. For 1 liter mixtures we need about 40,000 U Taq polymerase, 100 ml 2 mM NTP (100 mg NTP), 20 ml 100 μM DOP 6 M-W (2 μmol).

Example 4

The present invention includes an apparatus for doing large scale PCR preparations. The apparatus comprises a reaction tube, at least one pump for continuously

20 supplying of reagents to the reaction tube, four temperature chambers, and a collection vessel. The reagents are combined and enter the reaction tube by action of the pump. In some embodiments, there are two supply pumps which combine various components into a mixing vessel which then communicates with the reaction tube. The reaction solution flows through the reaction tube through a series of reaction tube lengths which alternately

25 pass through each of the four temperature chambers to produce a cycle segment. That is, the reaction solution flows through the reaction tube which passes through each of the four temperature chambers. The portion of the reaction tube which passes through each of the four temperature chambers is referred to as a cycle segment and the flow of the reaction solution through a cycle segment corresponds to a cycle. The reaction tube comprises at

least twenty consecutive cycle segments. That is the reaction tube emerges from the fourth chamber and reenters the first chamber at least twenty times. In preferred embodiments, the reaction tube comprises at least thirty consecutive cycle segments, in some preferred embodiments, the reaction tube comprises at least forty consecutive cycle segments.

5 In a preferred embodiment the genomic DNA is manufactured as copies using random PCR with partially degenerate primers. As discussed above, in some embodiments such as for example those involving severe genomic diseases, the total dosage for a course of treatment (for a few months) may comprise 10-20 g of DNA. For manufacturing of such amounts of DNA special large scale PCR apparatus is designed and
10 illustrated in Figure 4.

Appropriate proportions of Solution 1: PCR buffer, nucleoside triphosphates and DNA template in optional Reagent Vessel I (1) and Solution 2: Taq polymerase in optional Reagent Vessel II (2) are preferably first combined using two pumps, Pump I (3) and Pump II (4) and an optional mixing chamber (5) for gradient formation in HPLC. In some
15 embodiments, the mixed solutions are injection into the reaction tubing (6) by way of a single pump (not shown) which may be provided for example after a mixing chamber or, as shown here, optionally using two pumps (3, 4) where the DNA template is amplified in a reactor comprising four thermoregulated circulating water baths (7, 8, 9 and 10). The tubing (6) enters and exits all the four baths (7, 8, 9 and 10) one after another in series
20 which is referred to herein as a cycle. The temperature in the first bath (7) is preferably 95° C. The temperature in the second bath (8), which is referred to as the Cooling Bath I, is preferably 90° C. The temperature in the third bath (9), which is referred to as the Annealing Bath, is preferably 55° C. The temperature in the fourth bath (10) is preferably 72° C. The reaction mixture repeats the cycle for appropriate number of times (typically
25 30) which is specified in the optimization procedure. The cycle is repeated by the tubing (6) re-entering the bath (7) after emerging from bath (10) then continuing through bath (8), then bath (9), then (10) again and, to repeat back through the temperature chambers again (7, 8, 9 and 10).

The tubing can be made from any material compatible with PCR or its inner
30 surface can be coated with such material. The preferred material is stainless steel or plastic used for medical purposes. The inner diameter of the tubing is between 3 mm and 20 mm,

09753892.010301

but can also be wider. In some cases when tubing with inner diameter over 4 mm is used a rigid spiral is inserted in the tubing to provide automatic mixing of solutions inside the tubing. Optional Pump III (11) is used for injecting additional quantities of the reagents (NTPs, Taq polymerase) from optional Reagent Vessel III (12) into the main tubing. After
5 the reaction mixture has circulated through the last bath in the last cycle, it optionally moves through Cooling Bath II (12) and further into the tank for storage of the PCR product, the Collection Vessel (13). The temperature in the Cooling Bath II (12) and Collection Vessel (13) is preferably 0° C.

The machine is capable of performing PCR in 50-1000 liter of the reaction volume
10 per day. The apparatus provides for continuous flow of reaction mixture through the various baths. Depending upon the flow speed and length of tube within a given temperature bath, the length of time the reaction mixture will spend at a temperature during a cycle can be determined. The reaction times for each temperature is thus a function of the flow rate and the length of tube within a temperature bath. Various
15 configurations may be employed to achieve the desired lengths within the various temperature baths.

09753892-010301